



# Soils Map



State: **Iowa**  
 County: **Jefferson**  
 Location: **14-71N-8W**  
 Township: **Round Prairie**  
 Acres: **34.96**  
 Date: **12/6/2022**



Soils data provided by USDA and NRCS.

©2022 AgriData, Inc.

Area Symbol: IA101, Soil Area Version: 30

Code	Soil Description	Acres	Percent of field	CSR2 Legend	Non-Irr Class *c	*i Corn Bu	*i Soybeans Bu	CSR2**	CSR	*n NCCPI Soybeans
425D2	Keswick clay loam, 9 to 14 percent slopes, moderately eroded	10.19	29.1%		IVe	88.0	25.5	8	12	38
65E2	Lindley loam, 14 to 18 percent slopes, moderately eroded	8.79	25.1%		Vle	134.4	39.0	29	28	53
65F2	Lindley loam, 18 to 25 percent slopes, moderately eroded	6.32	18.1%		Vlle	110.4	32.0	10	8	38
730B	Nodaway-Coppock-Cantril complex, 2 to 5 percent slopes	3.35	9.6%		IIw	208.0	60.3	74	61	71
832C2	Weller silty clay loam, terrace, 5 to 9 percent slopes, moderately eroded	2.31	6.6%		IVe	80.0	23.2	61	40	69
S795D2	Ashgrove silty clay loam, 9 to 14 percent slopes, moderately eroded	1.39	4.0%		IVe	0.0	0.0	7		39
S132C2	Weller silty clay loam, 5 to 9 percent slopes, moderately eroded	1.25	3.6%		IIIe	0.0	0.0	59		67
1715	Nodaway-Vesser-Ackmore complex, 0 to 2 percent slopes	1.16	3.3%		IIw	190.4	55.2	74	80	83
132D2	Weller silty clay loam, 9 to 14 percent slopes, moderately eroded	0.20	0.6%		IVe	80.0	23.2	34	28	63
<b>Weighted Average</b>						<b>4.75</b>	<b>111.4</b>	<b>32.3</b>	<b>27.6</b>	<b>*n 49.7</b>

\*\*IA has updated the CSR values for each county to CSR2.

\*- CSR weighted average cannot be calculated on the current soils data, use prior data version for csr values.

\*i Yield data provided by the ISPAID Database version 8.1.1 developed by IA State University.

\*n: The aggregation method is "Weighted Average using all components"

\*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.