



# Soils Map



State: **Iowa**  
 County: **Hardin**  
 Location: **8-88N-22W**  
 Township: **Buckeye**  
 Acres: **147.4**  
 Date: **1/2/2025**



Maps Provided By:  
  
 CUSTOMIZED ONLINE MAPPING  
 © AgriData, Inc. 2023 www.AgriDataInc.com



Soils data provided by USDA and NRCS.

Area Symbol: IA083, Soil Area Version: 31

| Code                    | Soil Description   | Acres | Percent of field | CSR2 Legend | Non-Irr Class *c | *i Corn Bu  | *i Soybeans Bu | CSR2**      | CSR         | *n NCCPI Soybeans |
|-------------------------|--|-------|------------------|-------------|------------------|-------------|----------------|-------------|-------------|-------------------|
| L956                    | Harps-Okoboji complex, Bemis moraine, 0 to 2 percent slopes    | 51.01 | 34.6%            |             | Ilw              | 0.0         | 0.0            | 69          |             | 78                |
| L138B                   | Clarion loam, Bemis moraine, 2 to 6 percent slopes             | 30.95 | 21.0%            |             | Ile              | 220.8       | 64.0           | 88          |             | 79                |
| L507                    | Canisteo clay loam, Bemis moraine, 0 to 2 percent slopes       | 16.99 | 11.5%            |             | Ilw              | 0.0         | 0.0            | 87          |             | 81                |
| L95                     | Harps clay loam, Bemis moraine, 0 to 2 percent slopes          | 15.65 | 10.6%            |             | Ilw              | 0.0         | 0.0            | 75          |             | 80                |
| L107                    | Webster clay loam, Bemis moraine, 0 to 2 percent slopes        | 14.35 | 9.7%             |             | Ilw              | 0.0         | 0.0            | 88          |             | 83                |
| L55                     | Nicollet loam, 1 to 3 percent slopes                           | 13.14 | 8.9%             |             | Ie               | 0.0         | 0.0            | 91          |             | 84                |
| 6                       | Okoboji silty clay loam, 0 to 1 percent slopes                 | 4.12  | 2.8%             |             | Illw             | 185.6       | 53.8           | 59          | 59          | 74                |
| L329                    | Webster-Nicollet complex, Bemis moraine, 0 to 3 percent slopes | 0.82  | 0.6%             |             | Ilw              | 0.0         | 0.0            | 89          |             | 83                |
| 201B                    | Coland-Terril complex, 2 to 5 percent slopes                   | 0.37  | 0.3%             |             | Ilw              | 208.0       | 60.3           | 80          | 69          | 90                |
| <b>Weighted Average</b> |  |       |                  |             |                  | <b>1.94</b> | <b>52.1</b>    | <b>15.1</b> | <b>79.4</b> | <b>*n 79.7</b>    |

\*\*IA has updated the CSR values for each county to CSR2.

\*- CSR weighted average cannot be calculated on the current soils data, use prior data version for csr values.\*i Yield data provided by the ISPAID Database version 8.1.1 developed by IA State University.

\*n: The aggregation method is "Weighted Average using all components"

\*c: Using Capabilities Class Dominant Condition Aggregation Method